

# An Example of the $\Phi$ Function in a Biological System

James Lynn

## Abstract

This paper provides a material, numerical example of how the  $\Phi$  function can be used to evaluate the evolutionary efficiency of a simple biological system. We apply the  $\Phi$  function to a bacterial cell adapting to thermal stress, using real units for complexity, entropy, and energy efficiency. The results demonstrate how  $\Phi$  can be interpreted to assess the evolutionary dynamics of physical systems.

## 1. Introduction

The function is defined as a scalar that describes how a system evolves over time in terms of its internal complexity ( $C$ ), entropy ( $S$ ), and energy efficiency ( $E$ ). The general form is:

$$\Phi = \alpha \left( \frac{dC}{dt} \right) - \beta \left( \frac{dS}{dt} \right) + \gamma \left( \frac{dE}{dt} \right)$$

Where  $\alpha$ ,  $\beta$ , and  $\gamma$  are weights or scaling constants that normalize units and assign significance to each term.

## 2. Scenario: Bacterial Cell Responding to Heat

Consider a bacterial cell exposed to elevated temperature. In response:

- Internal regulatory complexity increases from 50 to 65 bits.
- Thermodynamic entropy rises from 2.0 to 2.5 J/K.
- Energy yield (ATP/glucose) increases from 32 to 34.

These changes occur over one hour, or 3600 seconds.

## 3. Step 1: Compute Rates of Change

$$\begin{aligned}\frac{dC}{dt} &= \frac{65 - 50}{3600} = 0.00417 \text{ bits/sec} \\ \frac{dS}{dt} &= \frac{2.5 - 2.0}{3600} = 0.000139 \text{ J/K/sec} \\ \frac{dE}{dt} &= \frac{34 - 32}{3600} = 0.000556 \text{ ATP/glucose/sec}\end{aligned}$$

## 4. Step 2: Define Scaling Coefficients

$$\alpha = 10\text{-units} \cdot \text{s/bit}, \quad \beta = 1000\text{-units} \cdot \text{s} \cdot \text{K/J}, \quad \gamma = 5\text{-units} \cdot \text{s/glucose-ATP}$$

## 5. Step 3: Calculate $\Phi$

$$\begin{aligned}\Phi &= 10 \cdot 0.00417 - 1000 \cdot 0.000139 + 5 \cdot 0.000556 \\ &= 0.0417 - 0.139 + 0.00278 \\ &= -0.0945 \Phi\text{-units}\end{aligned}$$

## 6. Interpretation

The result is:

$\Phi = -0.0945 \Phi\text{-units}$

This indicates that although complexity and energy efficiency increased, the rise in entropy outweighed these gains. The system is in a transitional stress phase and not yet optimizing its adaptive response.

## 7. Comparison: With and Without Using $\Phi$

- **Without  $\Phi$ :** Complexity and energy appear to improve, but entropy also increases. Its unclear how to weigh these effects together.
- **With  $\Phi$ :** The combined outcome is quantified as a single value. The negative  $\Phi$  value tells us the system is expending effort without a net gain in evolutionary quality.

| Feature           | Interpretation Without $\Phi$ | Rate or Value             | With $\Phi$ Weight                                                                       |
|-------------------|-------------------------------|---------------------------|------------------------------------------------------------------------------------------|
| Complexity        | Improves                      | +0.00417 bits/sec         | +0.0417                                                                                  |
| Entropy           | Increases                     | +0.000139 J/K/sec         | -0.139                                                                                   |
| Energy Efficiency | Slight gain                   | +0.000556 ATP/glucose/sec | +0.00278                                                                                 |
| <b>Net Effect</b> | Unclear/Mixed                 |                           | <div style="border: 1px solid black; padding: 2px; display: inline-block;">-0.0945</div> |

## 8. Conclusion

This example demonstrates how the  $\Phi$  function can be applied quantitatively using real units. It offers insight into the evolutionary trajectory of systems under pressure, providing a tool to measure adaptive success or failure using principles grounded in thermodynamics and information theory.